

^NFENTORATM

Fentanyl Buccal/Sublingual Effervescent Tablets
100 mcg, 200 mcg, 400 mcg, 600 mcg, and 800 mcg fentanyl as fentanyl citrate

Opioid Analgesic

Guide for Patients and Caregivers

Manufactured for:
Teva Canada Innovation
Montréal, Quebec H2Z 1S8

Distributed by:
Teva Canada Limited
Toronto, Ontario M1B 2K9

Guide for Patients and Caregivers

IMPORTANT DRUG INFORMATION

- FENTORA is a prescription opioid (narcotic) pain medicine containing fentanyl.
- FENTORA manages breakthrough pain in cancer patients who are already taking other opioid pain medicines continuously for cancer pain.
- FENTORA is only to be used by patients with cancer who are already taking opioid (narcotic) pain medicines every day and are used to them (opioid tolerant).

IMPORTANT SAFETY INFORMATION

- You should never give anyone your FENTORA. They could die from taking it. If a person has not been prescribed FENTORA, taking even one dose can cause a fatal overdose.
- Keep FENTORA in a safe place away from children.
- Accidental use by a child is a medical emergency and can result in death. Get emergency help right away if a child takes FENTORA by accident. If possible, try to remove FENTORA from the child's mouth.
- Taking FENTORA with other opioid medicines, benzodiazepines, alcohol, or other central nervous system depressants (including street drugs) can cause severe drowsiness, decreased awareness, breathing problems, coma, and death

For other important safety information, see the FENTORA Patient Package Insert: *Serious Warnings and Precautions*

Guide for Patients and Caregivers

THE MOST IMPORTANT INFORMATION YOU SHOULD KNOW ABOUT USING FENTORA

This guide is a summary and will not tell you everything about this drug. Talk to your healthcare professional about your medical condition and treatment and ask if there is any new information about **FENTORA**.

FENTORA, which contains the drug fentanyl, is a tablet that you place between your gum and cheek or under the tongue.

Fentanyl is a very strong opioid narcotic pain medicine that can cause serious and life-threatening breathing problems in the case of an overdose or if the dose you are using is too high for you. Get emergency medical help immediately if you:

- have trouble breathing, or have slow or shallow breathing
- have a slow heartbeat
- have extreme sleepiness
- have cold, clammy skin
- feel faint, dizzy, confused, or cannot think, walk, or talk normally
- have a seizure have hallucinations

If you have a troublesome symptom or side effect that is not listed here or becomes bad enough to interfere with your daily activities, talk to your healthcare professional.

Guide for Patients and Caregivers

GENERAL INFORMATION

Your doctor prescribed FENTORA for you to treat breakthrough cancer pain. Using FENTORA has some risks because it contains fentanyl, an opioid (narcotic) pain medicine.

FENTORA can cause life-threatening breathing problems if mistakenly taken by a child or by people who are not used to taking opioid pain medicines every day.

This booklet will help you learn more about FENTORA:

- Who should and should not use it
- How to correctly place FENTORA in your mouth
- How to dispose of unused units properly

You can get more information by consulting:

- A website that includes information about FENTORA and how to use it.
- A toll-free number [1-855-513-8382] to call if you have questions about FENTORA.
- A Patient Package Insert that gives important safety information.
- Color-coded packaging that helps you and your pharmacist make sure you are getting the strength of FENTORA your doctor prescribed.



Read this booklet carefully before starting FENTORA. Also, read the Patient Package Insert that comes with each prescription for FENTORA as it may contain updated information.

Ask the people who live with you to read this information. This booklet and the Patient Package Insert do not take the place of talking with your doctor about your breakthrough cancer pain. Ask your doctor or pharmacist any questions you have. Talk with them about any concerns you have.

Guide for Patients and Caregivers

FENTORA is for You and Only You

- Never give FENTORA to anyone else, even if they have the same symptoms you have. It may harm them and even cause death.
- Keep FENTORA in a safe place away from children and from anyone else for whom it has not been prescribed.

Use FENTORA Exactly as Prescribed

- FENTORA must be used only for the management of breakthrough cancer pain.
- Use FENTORA only if you are already taking other opioid pain medicines for your cancer pain every day and your body is used to these medicines (opioid tolerant). Your doctor will know if you are opioid tolerant. **Do not use FENTORA if you are not opioid tolerant.**
- You start taking FENTORA at the lowest dose of 100 micrograms (mcg). The dose goes up until you and your doctor find the right dose for you.
- Once your right dose has been selected, do not change your dose of FENTORA yourself. If you think you need more pain relief, talk with your doctor. Your doctor can decide if your dose of FENTORA needs to be changed.
- Do not take FENTORA more often than your doctor tells you.

Store FENTORA in a Safe, Secure Place

- Keep FENTORA in a safe place away from children.
- Accidental use by a child is a medical emergency and can result in death.
- Get emergency help right away if a child takes FENTORA by accident. If possible, try to remove FENTORA from the child's mouth.
- FENTORA is a controlled drug. It can be abused by people who abuse prescription medicines or street drugs.
- Keep FENTORA in a place where no one can easily reach it or steal it.
- FENTORA comes in a foil package. Do not open the package until ready to use. Once opened, use FENTORA right away.
- Follow the FENTORA administration guide (see below). Do not chew or swallow FENTORA. If you do, you will likely get less relief for your breakthrough pain.
- **Do not cut or split the FENTORA unit.**

Guide for Patients and Caregivers

FENTORA Administration Guide

Opening the Blister Package

1. Do not open the blister until ready to administer FENTORA.
2. Separate a single blister unit from the blister card by bending and tearing apart at the perforations.
3. Bend the blister unit along the line where indicated.
4. Peel back the blister backing to expose the tablet. **DO NOT attempt to push the tablet through the blister as this may cause damage to the tablet.**
5. Do not store the tablet once it has been removed from the blister package as this spoils the tablet and, more importantly, because this increases the risk of accidental exposure to the tablet.

Tablet Administration

- Once the tablet is removed from the blister unit, you should **immediately** place the entire FENTORA tablet in the buccal cavity (above a rear molar, between the upper cheek and gum) or under your tongue. **You should not split the tablet.**
- The FENTORA tablet should not be sucked, chewed or swallowed, as this will result in lower pain relief.
- The FENTORA tablet should be left between the cheek and gum, or under the tongue, until it has disintegrated, which usually takes approximately 14-25 minutes. After 30 minutes, if remnants from the FENTORA tablet remain, they may be swallowed with a glass of water.
- It is recommended that you switch sides of the mouth when administering subsequent doses of FENTORA.

FENTORA Dosing Instructions

To assist your doctor to find the right dose of FENTORA for you, and avoid the risk of overdosing, follow all instructions carefully.

- **All patients MUST begin treatment using 100 mcg FENTORA.**
- You will be closely followed by your doctor and your dosage strength will be changed until you reach the right dose for you
- **FENTORA should only be used ONCE for a breakthrough cancer pain episode. FENTORA should not be used again during the same episode.**
- During any episode of breakthrough cancer pain, if adequate pain relief *is not achieved* after FENTORA, you may use a rescue medication (other than FENTORA, after 30 minutes) *as directed by your doctor.*
- **You MUST wait at least 4 hours before treating another episode of breakthrough pain with FENTORA.**

Guide for Patients and Caregivers

- **Use of FENTORA should be limited to FOUR episodes of breakthrough pain per day. If you experience more than four breakthrough pain episodes per day, talk to your doctor.**

All unused tablets need to be returned to the pharmacy for proper disposal.

In case of drug overdose, contact a health care practitioner, hospital emergency department or regional Poison Control Centre immediately, even if there are no symptoms.

Follow FENTORA Dosing Instructions Carefully

- **FENTORA can cause life-threatening breathing problems which can lead to death.**
- **Call your doctor or get emergency medical help right away if you have any of the following symptoms:**
 - **Trouble breathing**
 - **Extreme sleepiness or drowsiness with slowed breathing**
 - **Slow, shallow breathing (little chest movement while breathing)**
 - **Faintness, dizziness, confusion, or other unusual symptoms**
 - **Inability to think, talk or walk normally**
 - **Seizure and hallucination**
 - **Abdominal pain, severe constipation, nausea**
 - **Agitation or restlessness, loss of muscle control or muscle twitching, tremor, diarrhea**
 - **Rash, hives, swelling of the face, lips, tongue or throat, difficulty swallowing or breathing (Allergic Reaction)**

Guide for Patients and Caregivers

ANSWERS TO COMMON QUESTIONS ABOUT FENTORA

Why has my doctor prescribed FENTORA?

Your doctor has prescribed FENTORA to treat your breakthrough cancer pain. The pain medicine you take every day on a regular basis helps control your constant pain. But you have also been feeling “flares” of intense pain that “breakthrough” the relief offered by your regular pain medicine. These flares are called “breakthrough pain” or BTP.

FENTORA is used specifically to treat BTP in patients with cancer who are already taking opioid (narcotic) pain medicine every day and are used to them (opioid tolerant).

What is FENTORA?

FENTORA delivers fentanyl in a tablet that you place between your gum and cheek



or under the tongue to deliver relief of breakthrough cancer pain.



Guide for Patients and Caregivers

What should I tell my doctor before I start taking FENTORA?

Tell your doctor about all your medical and mental health problems. Be sure to include:

- Trouble breathing or lung problems such as asthma, wheezing or being short of breath
- Head injury or brain problem
- Liver or kidney problems
- Seizures (convulsions or fits)
- Slow heart rate or other heart problems
- Low blood pressure
- Mental problems including major depression, schizophrenia or hallucinations (seeing or hearing things that are not there)
- Past or present problems with alcohol for you or a family member
- Past or present problems with drug abuse or addiction, or a family history of a drug abuse problem or addiction problem
- Migraines
- Chronic or severe constipation
- Problems with your thyroid, adrenal or prostate gland
- have any other medical conditions

Tell your doctor if you are:

- **Pregnant or planning to become pregnant.** FENTORA may harm your unborn baby.
- **Breastfeeding.** FENTORA passes through your breast milk and it can cause serious harm to your baby. You should not use FENTORA while breastfeeding.

Tell your doctor about all the medicines you take.

Include non-prescription medicines, vitamins, and herbal supplements. Some medicines may cause serious or life-threatening medical problems when taken with FENTORA.

How will my doctor know which FENTORA strength is right for me?

Your doctor or nurse will explain the step-by-step dosing instructions. These steps can help find out which strength of FENTORA is right for you.

How often should I use FENTORA?

Use FENTORA exactly as your doctor tells you. Do not take FENTORA more often than **4 times** a day. Separate each dose by at least **4 hours**.

Do I still use my regular medicine for persistent pain?

Keep using your regular pain medicine, unless your doctor tells you to stop it. Your regular pain medicine is long acting. It controls your constant pain. FENTORA does not take the place of your regular pain medicine. Use FENTORA only for breakthrough pain.

Guide for Patients and Caregivers

Who should NOT take FENTORA?

- People who are NOT already taking other opioid pain medicines every day for constant cancer pain.
- People who have severe respiratory problems
- People who are allergic to anything in FENTORA. The active ingredient is fentanyl.
- People who are currently taking monoamine-oxidase (MAO) inhibitors (used for severe depression) or have done so in the past 2 weeks.
- People who have any heart problems
- People who have bowel blockage or narrowing of the stomach or intestines
- People who have severe pain in their abdomen
- People who have a head injury
- People who are at risk for seizures
- People who suffer from alcoholism
- People who are going to have a planned surgery
- Women who are in labour
- Women who are breastfeeding

See the enclosed Patient Package Insert for a complete list of ingredients of FENTORA.

How will my pharmacist know which FENTORA strength to give me?

FENTORA units come in 5 different color-coded boxes for the 5 different dosage strengths. The color-coded boxes help your pharmacist give you the strength of FENTORA your doctor prescribed. The units are in color-coded child-resistant foil packages. Every unit is marked with a number to identify the dose it delivers.

Dosage Strength	Engraving on the Tablet	Carton/Blister Package Colour
100 mcg	1	Blue
200 mcg	2	Orange
400 mcg	4	Sage green
600 mcg	6	Magenta (pink)
800 mcg	8	Yellow

Can I eat or drink liquids while using FENTORA?

You can drink liquids after FENTORA has been in your mouth for 30 minutes but do not move the tablet(s) with your tongue/finger or eat any food until the tablet(s) has completely dissolved.

What happens if I bite, chew or swallow the FENTORA unit?

Do not bite, chew or swallow FENTORA units. If you do, you will likely get less relief for your breakthrough pain.

Guide for Patients and Caregivers

What are the most common side effects of FENTORA?

The most common side effects of FENTORA are

- nausea,
- vomiting,
- dizziness, fainting
- low red blood cell count,
- sleepiness,
- swelling of the arms, hands, legs and feet,
- headache
- constipation
- problems with vision
- itching
- sweating

FENTORA has other side effects. For a complete list, ask your doctor.

Talk with your doctor about any side effects that bother you or that do not go away.

Will FENTORA make me sleepy?

FENTORA can make you sleepy. Do not drive, operate machinery, or do other dangerous activities until you know how FENTORA affects you. Ask your doctor when it is okay to do those activities.

Can I become addicted to FENTORA?

FENTORA is an opioid medication. There is a chance of abuse or addiction with FENTORA.

If you are concerned, talk to your doctor or healthcare provider.

The chance of addiction is higher if you have ever been addicted to or abused other medications, street drugs, or alcohol, or have a history of mental health problems.

Can I take other medicines while I am using FENTORA?

- Tell your doctor about all your medicines. Include non-prescription medicines, vitamins, and herbal supplements.
- Some medicines may cause serious or life-threatening medical problems when taken with FENTORA
- The dose of certain medicines and FENTORA may need to be changed if you use them together.
- Do not start any new medicines while using FENTORA until you talk with your doctor. Your doctor will tell you if it is safe to take other medicines while you are using FENTORA.
- Be very careful about taking other medicines that make you sleepy, such as other pain

Guide for Patients and Caregivers

medicines or medicines for depression, sleep, nervousness, or allergies.

Can I drink alcohol while I am using FENTORA?

Do not drink alcohol while using FENTORA. It can increase your chance of having dangerous side effects.

What if a child or adult accidentally takes FENTORA?

If a child or adult accidentally takes FENTORA, call 911 right away for emergency help. If possible, try to immediately remove the FENTORA tablet from the mouth.

Where can I get more information on FENTORA?

If you have any further questions on the use of FENTORA, ask your doctor or pharmacist, or call Teva Canada Innovation for more information at the following telephone number:

1 855 513-8382

REFERENCE

1. FENTORA Canadian Product Monograph. Teva Canada Innovation, 07-OCT-2016.